

Clinical, Functional and Radiological Outcome of Posterior Lumbar Interbody Fusion by Banana Cage with Bone Graft for the Treatment of High-grade Lumbar Spondylolisthesis at L5-S1

Md. Anowarul Islam¹, Md. Mohoshin Sarker¹, Suvaradev Saha¹, Afia Ibnat Islam²

Abstract

Objectives: To evaluate the clinical, functional and radiological outcome of posterior lumbar interbody fusion (PLIF) by Banana cage with bone graft.

Material and Methods: This retrospective analytical study was carried out in spine unit of Orthopaedic surgery department of BSMMU, and other private hospital in Dhaka from January 2010 to December 2020. We did PLIF by Banana cage with bone graft for High-grade Lumbar Spondylolisthesis at L5-S1. The follow-up period ranges from 1- 2 years (average 18 months. All patients were assessed by VAS, ODI, WDI, Spino-pelvic parameters, Modified Macnab's Criteria to find out overall outcome and Hackenberge criteria for radiological fusion.

Results: Total 40 patients were included among 16 were male and 24 were female. The average age of the patients was 52.45 ± 10.1 years. 60.0% patients were housewife, 20.0%, 10.0%, 10% were day labourer, farmer and service holder respectively. Average pelvic tilt was $26.05 \pm 6.27^\circ$ preoperatively and $24.10 \pm 6.26^\circ$ at the final follow-up, average PI was $66.07 \pm 7.39^\circ$ preoperatively and $61.19 \pm 7.08^\circ$ at the final follow-up. Preoperative lumbar lordosis was $45.55 \pm 6.71^\circ$ & at final follow-up $37.29 \pm 6.19^\circ$. VAS score and ODI scales were improved significantly from preoperative 6.90 ± 6.16 and 57.60 ± 15.66 to at final follow-up 2.0 ± 0.8 and 7.60 ± 2.40 . Pre-operative Translation ratio, slip angle and disc height ratio were 21.96 ± 10.25 , -18.87 ± 8.28 , 11.03 ± 4.36 and at postoperatively 13.17 ± 6.57 , -18.44 ± 7.12 , 19.60 ± 3.36 . Fusion was achieved in 36 cases (90%), 3 cases (7.5%) were fragmented and pseudoarthrosis showed only 1 case (2.5%). Post-operative clinical outcome showed excellent outcome (95%), 1(2.5%) case had good and 1(2.5%) case had fair outcome.

Conclusion: It can be concluded that, PLIF by Banana cage with bone graft is a very good option for the treatment of High-grade Lumbar Spondylolisthesis at L5-S1 levels.

Keywords: PLIF, High-grade Lumbar Spondylolisthesis, Pelvic parameters, Banana Cage.

Introduction

Pain in the lower lumbar region is a socioeconomically serious medical illness worldwide. The main reason from physiological consideration is micro and macro instability of spine [1, 2]. There are numerous causes for backache. Spondylolisthesis is one among them [3]. Spondylolisthesis is translation of one vertebra over the next lower vertebra. High-

grade spondylolisthesis (HGS) is defined as more than 50% slippage of a vertebral body relative to an adjacent one, as per Meyerding classification, and most common location being L5/S1 followed by L4/L5 [4]. Most commonly used classification systems for spondylolisthesis were introduced by Wiltse et al [5] and Marchetti and Bartolozzi [6] which is most practical classification system in terms of prognosis and therapy.

In addition to the bony morphologic changes seen in high-dysplastic spondylolisthesis, spinopelvic balance plays an important role in developing and progression of spondylolisthesis [7]. Altered biomechanical stresses found due to abnormal spino-pelvic balance at the lumbosacral junction and compensatory mechanisms used to maintain adequate posture and gait.



¹Department of Spine Surgery, Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh.

²Department of Spine Surgery, Holy Family & Red Crescent Medical College & Hospital, Dhaka, Bangladesh.

Address of correspondence :

Dr. Md. Anowarul Islam,
Professor & Unit Head of Spine Surgery, Bangabandhu
Sheikh Mujib Medical University, Dhaka, Bangladesh.

E-mail: maislam.spine@gmail.com

Submitted: 21/03/2022; Reviewed: 19/04/2022; Accepted: 17/07/2022; Published: 01/10/2022

Back Bone: The Spine Journal (The Official Journal Of "Spine Association of Gujarat") | Available on www.backbonejournal.com | DOI:10.13107/bbj.2022.v03i02.046

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial-Share Alike 4.0 License (<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which allows others to remix, tweak, and build upon the work non-commercially as long as appropriate credit is given and the new creation are licensed under the identical terms.

Degenerative lumbar spondylolisthesis (DLS) is always associated with facet joint degeneration and mostly observed in persons over the age of 50 years. Patient of spinal stenosis may suffer from back and/or leg pain [8]. Decompression alone may further increase the instability, thereby decompression with stabilization is better [9, 10].

Vertebral Interbody fusion (IBF), is relatively new set of technique, has become very popular in the treatment of symptomatic DLS. Interbody fusion improves the biomechanical stability by stabilizing the anterior column, specially in patients with High-grade spondylolisthesis, Unstable slips, Degenerative type of scoliosis, and retained disc height etc. [11–14]. Insertion of interbody devices also improve sagittal alignment and restore disc and foraminal height as well, which ultimately provide indirect decompression of foraminal and canal stenosis and aiding in spondylolisthesis reduction [12, 15].

Posterior lumbar interbody fusion (PLIF) surgery is performed by the standard posterior approach. Wide laminectomy is done first followed by partial bilateral facetectomy, then the neural elements are retracted to either side, to make space for disc space preparation and finally insertion of a titanium interbody device packed with autogenous bone graft within the inter-vertebral space [16, 17].

Material and Methods

This retrospective study of 40 patients were carried out in the spine unit of Orthopaedic surgery department of Bangabandhu Sheikh Mujib medical University, and other private hospitals in Dhaka, Bangladesh from January 2010 to December 2020. PLIF by Banana cage with bone graft was done for High-grade Lumbar Spondylolisthesis only at L5-S1.




Figure 1: Sacral slope (SS) is angle subtended by horizontal reference line (HRL) and sacral endplate line (bc). SS shares common reference line (bc) with pelvic incidence (PI) and pelvic tilt (PT). PI is measured from static anatomic structures. PT and SS depend on angular position of sacrum/pelvis in relation to femoral heads, which changes with standing, sitting, and lying down. Relationship of PT and SS is affected by lumbosacropelvic flexion and extension. VRL, Vertical reference line. (From Jackson R, Kanemura T, Kawakami N, Hales C: Lumbopelvic lordosis and pelvic balance on repeated standing lateral radiographs of adult volunteers and untreated patients with constant low back pain, Spine 25:575–586, 2000.)

The follow up period ranges from 1 year to 2 years (average 18 months). Within these follow up period we have assessed the patients clinically, functionally and radiologically. All patients were assessed pre and post operatively by Visual Analogue Score (VAS), Oswestry Disability Index (ODI), Waddell disability index (WDI), Spino-pelvic parameters, Modified Macnab's Criteria to find out overall outcome and Hackenberge criteria for radiological fusion. IBM-SPSS V26 software was used for statistical analysis.

Selection of the patients for surgery

Inclusion criteria: Patients with grade iii, iv and v spondylolisthesis, only at L5-S1 who had severe low pain, neurological deficit and restriction of movement with instability not responding to conservative treatment.

Exclusion criteria: Patients with grade i and grade ii spondylolisthesis. High grade spondylolisthesis patients with severe comorbidities and infection at local incision site.

Operative technique:

Patient is prone and two parallel pillows on a radiolucent table after General anaesthesia ensure the abdomen hangs free. After proper cleaning, painting and draping of the operative area, a lower midline incision was made extending from spinous process of L4 to S3. The skin, subcutaneous tissue was cut in a single line, and the deep fascia and supra spinous ligaments cut in a same plane by diathermy. Then para spinal muscles made

0% to 20%: minimal disability:	The patient can cope with most living activities. Usually no treatment is indicated apart from advice on lifting sitting and exercise.
21%-40%: moderate disability:	The patient experiences more pain and difficulty with sitting, lifting and standing. Travel and social life are more difficult and they may be disabled from work. Personal care, sexual activity and sleeping are not grossly affected and the patient can usually be managed by conservative means.
41%-60%: severe disability	Pain remains the main problem in this group but activities of daily living are affected. These patients require a detailed investigation.
61%-80%: crippled:	Back pain impinges on all aspects of the patient's life. Positive intervention is required.
81%-100%:	These patients are either bed-bound or exaggerating their symptoms.

Chart 1: Oswestry Disability index

Parameters	Description
Sacral slope (SS)	The angle between the horizontal line and the cranial sacral endplate tangent
Pelvic tilt (PT)	The angle between the vertical line and the line joining the middle of the sacral plate to the centre of the bicoxfemoral axis
Pelvic index (PI)	The angle between the line perpendicular to the middle of the cranial sacral endplate and the line joining the middle of the cranial sacral endplate to the centre of the bicoxfemoral axis
Lumber lordosis (LL)	Cobb angle measured from the superior endplate of L1 to the superior endplate of S1
C7 Plumb line	Vertical line drawn from the centre of C7 vertebrae on a radiograph. Often used as a reference line for measuring sagittal balance. The distal reference point for this parameter is the posteriorsuperior corner of the sacrum

chart 2: Spino-pelvic parameters



Figure 2: Pre-operative X-ray



Figure 3: Pre-operative magnetic resonance imaging



Figure 4: Banana cage filled with bone graft, Per-operative picture showing pedicle screw and rod





Figure 5: Per-operative C-ARM imaging



Figure 6: immediate Post operative -operative X-ray

Figure 7: 18th month Post-operative X-ray showing good bony fusion

retracted subperiosteally up-to tip of transverse processes bilaterally. The space was identified under C-ARM guidance and partial or complete laminectomy of L5 was done. After identification of L5 nerve root, L5-S1 joint space clearance was done using end plate curette and box curette. After that we put pedicle screw from L4 to S1 on either side. Then distraction between L4 and S1 done after putting pre bended rod on the pedicle screws slots, reduction of L5 achieved by either using a persuator with a traditional L5 pedicle screw or sometimes by a reduction screw. After reduction we again distract between L5-

S1 to get enough space for introduction of proper sized Banana cage with bone graft. After putting cage with bone graft at space, we put a gel foam over open cauda equina. Proper haemostasis was ensured in every step of surgery. The wound was closed in anatomical layer by absorbable 1/0 cutting body vicryl and skin by skin stapler keeping a drain in situ.

Results

The retrospective analytical study includes 40 patients who fulfilled the inclusion criteria, were operated by PLIF by

Score	Pre operative (Mean ± SD)	Final follow-up (Mean ± SD)	p value
VAS	6.90 ± 6.16	2.0 ± 0.8	<0.001
WDI	7.03 ± 1.08	2.07 ± 0.61	<0.001

Pre operative and post operative comparison of clinical outcomes after 24 months (n=40)
According to VAS & WDI score

	Pre operative	24 months after operation	P value
Pelvic Tilt (%)	26.05 ± 6.27°	24.10 ± 6.26°	<0.001
Pelvic Index (°)	66.07 ± 7.39°	61.19 ± 7.08°	<0.001
Lumbar Lordosis (°)	45.55 ± 6.71°	37.29 ± 6.19°	<0.001

Mean pre- and post-operative (after 24 months) spinopelvic parameters (n=40)

Comprehensive outcome	Frequency (n)	Percentage (%)
Excellent	35	87
Good	4	10
Fair	1	2.5
Poor	0	0

Distribution of patients according to post operative clinical outcome (n=40)
According to Modified Macnab criteria (Macnab I et al.1971)

Banana cage with bone graft for High-grade Lumbar Spondylolisthesis. All patient were followed up from 1 year to 2 years (average 18 months) postoperatively.

Discussion

Patients treated by PLIF with Banana cage and bone graft were followed up for the period of 24 months. Overall clinical outcome categorized as excellent, good, fair, poor according to modified Macnab criteria. For statistical analysis good and excellent were grouped as satisfactory and fair and poor as unsatisfactory.

In this study, age range of the patients were 41–72 years, mean age was 52.45 ± 10.1 years and the male to female ratio was 8:12 which are comparable to the study of Audat et al, 2012. Who found mean age 54.2 ± 13.6 ; (range 36.0–66.0) [18].

Regarding occupation of the patient maximum (60.0%) patients were housewife followed by 20.0%, 10.0%, 10.0% were day labourer, farmer and service holder respectively. In study of Sakeb et al., 2013 maximum 65.4% were house wife, 15.4% were manual worker and 19.2% were sedentary worker which is also comparable [19].

All 40 (100%) patients had back pain, sciatica and neurogenic

Score	Pre operative (Mean ± SD)	Final follow-up (Mean ± SD)	p value
ODI	57.60 ± 15.66	7.60 ± 2.40	<0.001

Pre operative and post operative comparison of clinical outcomes after 24 months (n=40)
According to ODI score

24 months after operation n (%)	
Fused 36	90
Probably fused 3	7.5
Pseudo-arthrosis 1	2.5
Total 40 (100.0)	100

Radiological fusion status after 24 months of operation (n=40)
According to Hackenberge criteria (Hackenberge et al.2005)

No. of Patient develop fusion (n=40)	Time taken for fusion (Radiological Fusion by X-ray)				(Mean ± SD)
	6th month	12th month	18th month	18 month (CT scan)	
	22	7	5	5	
	-55%	-17.50%	-12.50%	-12.50%	9.9 ± 4.32

Distribution of patients according to time taken for fusion (n=40)

claudication (Audat et al., 2012). [18], Sensory disturbance was present in 38(95%) patients, and motor weakness was observed in 31(77.5%) patients. Only 1(2.5%) patient had urinary retention. Which is also comparable to study of Sakeb et al., 2013 [19]. All were improved significantly ($P<.005$) in postoperative group.

Decrease in spinopelvic parameters were observed after operation and remain almost unchanged throughout the follow up period (average pelvic tilt was $26.05 \pm 6.27^\circ$ preoperatively and $24.10 \pm 6.26^\circ$ at the final follow-up, average PI was $66.07 \pm 7.39^\circ$ preoperatively and $61.19 \pm 7.08^\circ$ at the final follow-up. Pre-operative lumbar lordosis were $45.55 \pm 6.71^\circ$ and $37.29 \pm 6.19^\circ$ at final follow-up). Study carried out by Lengert et al, 2014 [20]. Showed similar improvement.

Significant improvement of VAS score for back pain and post operative score from 6.90 ± 6.16 to 2.0 ± 0.8 score with significant ($p<0.001$), and ODI was significant before operation 57.60 ± 15.66 to 7.60 ± 2.40 with ($P<0.001$) in at 12 months follow up. In a study of Sakeb et al (2013) mean VAS reduced from 7.2 to 2.2 and mean ODI reduce from 60.7 to 11.2 at 12 months follow up. WDI significantly reduced from 7.03 ± 1.08 to 2.07 ± 0.61 with a significant ($P<0.001$) value

[19].

About 55% achieved radiological fusion at 6th month in our series which co-inside with 52% early fusion in Lee H et al. (2012) series [21]. Rate of spinal fusion with bone graft range from 46%-90% in Lowe GT et al. (2002) series [22]. In our series fusion evaluation by CT scan for 5 patient who showing doubt in x-ray fusion, 4 out of 5(80.00%) fused and one was fragmented according Cristensen assessment scale after 18 months of follow-up. Fusion after 18 months including X-ray & CT scan evaluation is 97.5%. Fusion was achieved 95.7% (45 of 47) cases in the study of AGAZZI et al. (1999) [23]. Radiographic fusion was present in 27(88.9%) patients after one year in the study of Audat et al. (2012) [18].

Post operative wound infection was in 1(2.5%) and pseudoarthrosis was in 1(2.5%) patient. Wound infection was managed conservatively by antibiotics according to culture and sensitivity report, improvement of nutritional status, removal of stitch, regular dressing secondary wound closure. Pseudoarthrosis was managed conservatively as the symptoms reduced significantly and patient continues his daily life without any trouble.

References

- Clinical spinal instability and low back pain, Manohar M. Panjabi, Journal of Electromyography and Kinesiology. 2003; 13:371-379.
- Lin Y, Chen W, Chen A, Li F. Comparison between minimally invasive and open transforaminal lumbar interbody fusion: A meta-analysis of clinical results and safety outcomes. J Neurol Surg A Cent Eur Neurosurg 2016;77:2-10.
- Karla R, Kumar NP, Kalyan SS. A study on management of high grade spondylolisthesis.
- Meyerding HW. Spondylolisthesis; surgical fusion of lumbosacral portion of spinal column and interarticular facets; use of autogenous bone grafts for relief of disabling backache. J Int Coll Surg., 1956; 26: 566-91.
- Wiltse LL, Newman PH, Macnab I (1976) Classification of spondylolysis and spondylolisthesis. Clin Orthop Relat Res 117:23-29
- Marchetti PG, Bartolozzi P (1997) Classification of spondylolisthesis as a guideline for treatment. In: Bridwell K, DeWald R (eds) The textbook of spinal surgery, 2nd edn edn. LippincottRaven, Philadelphia, pp 1211-1254
- Vidal J, Marnay T (1983) Morphology and anteroposterior body equilibrium in spondylolisthesis L5/S1. Rev Chir Orthop 69(1):17-28
- Jacobsen S, Sonne-Holm S, Rovsing H, Monrad H, Gebuhr P. Degenerative lumbar spondylolisthesis: an epidemiological perspective: the Copenhagen Osteoarthritis Study. Spine. 2007;32(1):
- Kepler CK, Vaccaro AR, Hilibrand AS, Anderson DG, Rihn JA, Albert TJ, et al. National trends in the use of fusion techniques to treat degenerative spondylolisthesis. Spine. 2014;39(19):1584-9.
- Schroeder GD, Kepler CK, Kurd MF, Vaccaro AR, Hsu WK, Patel AA, Savage JW. Rationale for the Surgical Treatment of Lumbar Degenerative Spondylolisthesis. Spine (Phila Pa 1976). 2015;40(21):E1161-6.
- Oda I, Abumi K, Yu B-S, Sudo H, Minami A. Types of spinal instability that require interbody support in posterior lumbar reconstruction: an in vitro biomechanical investigation. Spine. 2003;28(14):1573-80.
- McAfee PC, DeVine JG, Chaput CD, Prybis BG, Fedder IL, Cunningham BW, et al. The indications for interbody fusion cages in the treatment of spondylolisthesis: analysis of 120 cases. Spine. 2005;30(6 Suppl):S60-5.
- Liao J-C, Lu M-L, Niu C-C, Chen W-J, Chen L-H. Surgical outcomes of degenerative lumbar spondylolisthesis with anterior vacuum disc: can the intervertebral cage overcome intradiscal vacuum phenomenon and enhance posterolateral fusion? J Orthop Sci Off J Jpn Orthop Assoc. 2014;19(6):851-9.
- Ha K-Y, Na K-H, Shin J-H, Kim K-W. Comparison of posterolateral fusion with and without additional posterior lumbar interbody fusion for degenerative lumbar spondylolisthesis. J Spinal Disord Tech. 2008;21(4):229-34.
- Vamvanij V, Ferrara LA, Hai Y, Zhao J, Kolata R, Yuan HA. Quantitative changes in spinal canal dimensions using interbody distraction for spondylolisthesis. Spine. 2001;26(3):E13-8.
- Cole CD, McCall TD, Schmidt MH, Dailey AT. Comparison of lowback fusion techniques: transforaminal lumbar interbody fusion (TLIF) or posterior lumbar interbody fusion (PLIF) approaches. Curr Rev Musculoskelet Med. 2009;2(2):118-26.
- Lara-Almunia M, Gomez-Moreta JA, Hernandez-Vicente J. Posterior lumbar interbody fusion with instrumented posterolateral fusion in adult spondylolisthesis: description and association of clinico-surgical variables with prognosis in a series of 36 cases. Int J Spine Surg. 2015;9:22.
- Audat Z, Moutasem O, Yousef K, Mohammad B. Comparison of clinical and radiological results of posterolateral fusion, posterior lumbar interbody fusion and transforaminal lumbar interbody fusion techniques in the treatment of degenerative lumbar spine. Singapore medical journal. 2012 Mar 1;53(3):183-7.

In our study 35(87.5%) patients got satisfactory results and 4(10.0%) patients got good result in 1(2.5%) case had fair outcome.

Conclusion

The results of the our study indicate that the outcomes were significantly improved with surgery and the surgical technique of Posterior lumbar interbody fusion with local autogenous bone graft and Cage produced satisfying clinical, functional and radiological improvement.

19. Sakeb PN, Ahsan K. Comparison of the early results of transforaminal lumbar interbody fusion and posterior lumbar interbody fusion in symptomatic lumbar instability. Indian journal of orthopaedics. 2013 Jun;47(3):255-63.
20. Lengert R, Charles YP, Walter A, Schuller S, Godet J, Steib JP. Posterior surgery in high-grade spondylolisthesis. Orthopaedics & Traumatology: Surgery & Research. 2014 Sep 1;100(5):481-4.
21. Lee KH, Yue WM, Yeo W, Soeharno H, Tan SB. Clinical and radiological outcomes of open versus minimally invasive transforaminal lumbar interbody fusion. European spine journal. 2012 Nov;21(11):2265-70.
22. Lowe TG, Tahernia AD, O'Brien MF, Smith DA. Unilateral transforaminal posterior lumbar interbody fusion (TLIF): indications, technique, and 2-year results. Clinical Spine Surgery. 2002 Feb 1;15(1):31-8.
23. Agazzi S, Reverdin A, May D. Posterior lumbar interbody fusion with cages: an independent review of 71 cases. Journal of Neurosurgery: Spine. 1999 Oct 1;91(2):186-92.
24. Moreau S, Lonjon G, Guigui P, Lenoir T, de Loubresse CG, Chopin D. Reduction and fusion in high-grade L5-S1 spondylolisthesis by a single posterior approach. Results in 50 patients. Orthopaedics & Traumatology: Surgery & Research. 2016 Apr 1;102(2):233-7.
25. Ha KY, Na KH, Shin JH, Kim KW. Comparison of posterolateral fusion with and without additional posterior lumbar interbody fusion for degenerative lumbar spondylolisthesis. Clinical Spine Surgery. 2008 Jun 1;21(4):229-34.
26. Hackenberg L, Halm H, Bullmann V, Vieth V, Schneider M, Liljenqvist U. Transforaminal lumbar interbody fusion: a safe technique with satisfactory three to five year results. European Spine Journal. 2005 Aug;14(6):551-8.
27. Enker PA, Steffee AD. Interbody fusion and instrumentation. Clinical Orthopaedics and Related Research. 1994 Mar 1(300):90-101.
28. Meyers LL, Dobson SR, Wiegand D, Webb JD, Mencio GA. Mechanical instability as a cause of gait disturbance in high-grade spondylolisthesis: a pre-and postoperative three-dimensional gait analysis. Journal of Pediatric Orthopaedics. 1999 Sep 1;19(5):672.
29. Vaz G, Roussouly P, Berthonnaud E, Dimnet J. Sagittal morphology and equilibrium of pelvis and spine. European spine journal. 2002 Feb;11(1):80-7.
30. Vialle R, Charosky S, Padovani JP, Rigault P, Glorion C. Surgical treatment of high-grade lumbosacral spondylolisthesis in childhood, adolescent and young adult by the "double-plate" technique: a past experience. European Spine Journal. 2006 Aug;15(8):1210-8.

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given his/her consent for his/her images and other clinical information to be reported in the Journal. The patient understands that his/her name and initials will not be published, and due efforts will be made to conceal his/her identity, but anonymity cannot be guaranteed.

Conflict of Interest: NIL
Source of Support: NIL

How to Cite this Article

Islam MA, Sarker MM, Saha S, Islam AI | Clinical, Functional and Radiological Outcome of Posterior Lumbar Interbody Fusion by Banana Cage with Bone Graft for the Treatment of High-grade Lumbar Spondylolisthesis at L5-S1 | Back Bone: The Spine Journal | October 2022-March 2023; 3(2): 90-95.